


Part I

1. What is the x -coordinate of the point of inflection on the graph of $y = \frac{1}{3}x^3 + 5x^2 + 24$?

- (A) 5 (B) 0 (C) $-\frac{10}{3}$ (D) -5 (E) -10



2. The graph of a piecewise-linear function f , for $-1 \leq x \leq 4$, is shown above. What is the value of

$$\int_{-1}^4 f(x) dx ?$$

- (A) 1 (B) 2.5 (C) 4 (D) 5.5 (E) 8

3. $\int_1^2 \frac{1}{x^2} dx =$

- (A) $-\frac{1}{2}$ (B) $\frac{7}{24}$ (C) $\frac{1}{2}$ (D) 1 (E) $2 \ln 2$

4. If f is continuous for $a \leq x \leq b$ and differentiable for $a < x < b$, which of the following could be false?

(A) $f'(c) = \frac{f(b) - f(a)}{b - a}$ for some c such that $a < c < b$.

(B) $f'(c) = 0$ for some c such that $a < c < b$.

(C) f has a minimum value on $a \leq x \leq b$.

(D) f has a maximum value on $a \leq x \leq b$.

(E) $\int_a^b f(x) dx$ exists.

5. $\int_0^x \sin t dt =$

- (A) $\sin x$ (B) $-\cos x$ (C) $\cos x$ (D) $\cos x - 1$ (E) $1 - \cos x$

6. If $x^2 + xy = 10$, then when $x = 2$, $\frac{dy}{dx} =$

- (A) $-\frac{7}{2}$ (B) -2 (C) $\frac{2}{7}$ (D) $\frac{3}{2}$ (E) $\frac{7}{2}$

7. $\int_1^e \left(\frac{x^2 - 1}{x} \right) dx =$

- (A) $e - \frac{1}{e}$ (B) $e^2 - e$ (C) $\frac{e^2}{2} - e + \frac{1}{2}$ (D) $e^2 - 2$ (E) $\frac{e^2}{2} - \frac{3}{2}$


8. Let f and g be differentiable functions with the following properties:

(i) $g(x) > 0$ for all x

(ii) $f(0) = 1$

If $h(x) = f(x)g(x)$ and $h'(x) = f(x)g'(x)$, then $f(x) =$

- (A) $f'(x)$ (B) $g(x)$ (C) e^x (D) 0 (E) 1



9. The flow of oil, in barrels per hour, through a pipeline on July 9 is given by the graph shown above. Of the following, which best approximates the total number of barrels of oil that passed through the pipeline that day?

- (A) 500 (B) 600 (C) 2,400 (D) 3,000 (E) 4,800

10. What is the instantaneous rate of change at $x = 2$ of the function f given by $f(x) = \frac{x^2 - 2}{x - 1}$?

- (A) -2 (B) $\frac{1}{6}$ (C) $\frac{1}{2}$ (D) 2 (E) 6


11. If f is a linear function and $0 < a < b$, then $\int_a^b f''(x) dx =$

- (A) 0 (B) 1 (C) $\frac{ab}{2}$ (D) $b - a$ (E) $\frac{b^2 - a^2}{2}$


12. If $f(x) = \begin{cases} \ln x & \text{for } 0 < x \leq 2 \\ x^2 \ln 2 & \text{for } 2 < x \leq 4, \end{cases}$ then $\lim_{x \rightarrow 2} f(x)$ is

- (A) $\ln 2$ (B) $\ln 8$ (C) $\ln 16$ (D) 4 (E) nonexistent

APCalculus1998AB




13. The graph of the function f shown in the figure above has a vertical tangent at the point $(2, 0)$ and horizontal tangents at the points $(1, -1)$ and $(3, 1)$. For what values of x , $-2 < x < 4$, is f not differentiable?
- (A) 0 only (B) 0 and 2 only (C) 1 and 3 only (D) 0, 1, and 3 only (E) 0, 1, 2, and 3
14. A particle moves along the x -axis so that its position at time t is given by $x(t) = t^2 - 6t + 5$. For what value of t is the velocity of the particle zero?
- (A) 1 (B) 2 (C) 3 (D) 4 (E) 5
15. If $F(x) = \int_0^x \sqrt{t^3 + 1} dt$, then $F'(2) =$
- (A) -3 (B) -2 (C) 2 (D) 3 (E) 18
16. If $f(x) = \sin(e^{-x})$, then $f'(x) =$
- (A) $-\cos(e^{-x})$
 (B) $\cos(e^{-x}) + e^{-x}$
 (C) $\cos(e^{-x}) - e^{-x}$
 (D) $e^{-x} \cos(e^{-x})$
 (E) $-e^{-x} \cos(e^{-x})$








17. The graph of a twice-differentiable function f is shown in the figure above. Which of the following is true?
- (A) $f(1) < f'(1) < f''(1)$
 (B) $f(1) < f''(1) < f'(1)$
 (C) $f'(1) < f(1) < f''(1)$
 (D) $f''(1) < f(1) < f'(1)$
 (E) $f''(1) < f'(1) < f(1)$

18. An equation of the line tangent to the graph of $y = x + \cos x$ at the point $(0, 1)$ is
 (A) $y = 2x + 1$ (B) $y = x + 1$ (C) $y = x$ (D) $y = x - 1$ (E) $y = 0$
19. If $f''(x) = x(x + 1)(x - 2)^2$, then the graph of f has inflection points when $x =$
 (A) -1 only (B) 2 only (C) -1 and 0 only (D) -1 and 2 only (E) $-1, 0,$ and 2 only
20. What are all values of k for which $\int_{-3}^k x^2 dx = 0$?
 (A) -3 (B) 0 (C) 3 (D) -3 and 3 (E) $-3, 0,$ and 3
21. If $\frac{dy}{dt} = ky$ and k is a nonzero constant, then y could be
 (A) $2e^{kty}$ (B) $2e^{kt}$ (C) $e^{kt} + 3$ (D) $kt y + 5$ (E) $\frac{1}{2}ky^2 + \frac{1}{2}$
22. The function f is given by $f(x) = x^4 + x^2 - 2$. On which of the following intervals is f increasing?
 (A) $\left(-\frac{1}{\sqrt{2}}, \infty\right)$
 (B) $\left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$
 (C) $(0, \infty)$
 (D) $(-\infty, 0)$
 (E) $\left(-\infty, -\frac{1}{\sqrt{2}}\right)$



23. The graph of f is shown in the figure above. Which of the following could be the graph of the derivative of f ?






24. The maximum acceleration attained on the interval $0 \leq t \leq 3$ by the particle whose velocity is given by $v(t) = t^3 - 3t^2 + 12t + 4$ is
- (A) 9 (B) 12 (C) 14 (D) 21 (E) 40
25. What is the area of the region between the graphs of $y = x^2$ and $y = -x$ from $x = 0$ to $x = 2$?
- (A) $\frac{2}{3}$ (B) $\frac{8}{3}$ (C) 4 (D) $\frac{14}{3}$ (E) $\frac{16}{3}$


x	0	1	2
$f(x)$	1	k	2

26. The function f is continuous on the closed interval $[0, 2]$ and has values that are given in the table above. The equation $f(x) = \frac{1}{2}$ must have at least two solutions in the interval $[0, 2]$ if $k =$
- (A) 0 (B) $\frac{1}{2}$ (C) 1 (D) 2 (E) 3
27. What is the average value of $y = x^2\sqrt{x^3 + 1}$ on the interval $[0, 2]$?
- (A) $\frac{26}{9}$ (B) $\frac{52}{9}$ (C) $\frac{26}{3}$ (D) $\frac{52}{3}$ (E) 24
28. If $f(x) = \tan(2x)$, then $f'\left(\frac{\pi}{6}\right) =$
- (A) $\sqrt{3}$ (B) $2\sqrt{3}$ (C) 4 (D) $4\sqrt{3}$ (E) 8

1.D 2.B 3.C 4.B 5.E 6.A 7.E 8.E 9.D 10.D 11.A 12.E 13.B 14.C
 15.D 16.E 17.D 18.X 19.C 20.A 21.B 22.C 23.A 24.D 25.D 26.A
 27.A 28.E



76. The graph of a function f is shown above. Which of the following statements about f is false?
- (A) f is continuous at $x = a$.
- (B) f has a relative maximum at $x = a$.
- (C) $x = a$ is in the domain of f .
- (D) $\lim_{x \rightarrow a^+} f(x)$ is equal to $\lim_{x \rightarrow a^-} f(x)$.
- (E) $\lim_{x \rightarrow a} f(x)$ exists.
77. Let f be the function given by $f(x) = 3e^{2x}$ and let g be the function given by $g(x) = 6x^3$. At what value of x do the graphs of f and g have parallel tangent lines?
- (A) -0.701
- (B) -0.567
- (C) -0.391
- (D) -0.302
- (E) -0.258
78. The radius of a circle is decreasing at a constant rate of 0.1 centimeter per second. In terms of the circumference C , what is the rate of change of the area of the circle, in square centimeters per second?
- (A) $-(0.2)\pi C$
- (B) $-(0.1)C$
- (C) $-\frac{(0.1)C}{2\pi}$
- (D) $(0.1)^2 C$
- (E) $(0.1)^2 \pi C$
80. The first derivative of the function f is given by $f'(x) = \frac{\cos^2 x}{x} - \frac{1}{5}$. How many critical values does f have on the open interval $(0, 10)$?
- (A) One
- (B) Three
- (C) Four
- (D) Five
- (E) Seven



79. The graphs of the derivatives of the functions f , g , and h are shown above. Which of the functions f , g , or h have a relative maximum on the open interval $a < x < b$?

- (A) f only
 (B) g only
 (C) h only
 (D) f and g only
 (E) f , g , and h

80. The first derivative of the function f is given by $f'(x) = \frac{\cos^2 x}{x} - \frac{1}{5}$. How many critical values does f have on the open interval $(0, 10)$?

- (A) One
 (B) Three
 (C) Four
 (D) Five
 (E) Seven

81. Let f be the function given by $f(x) = |x|$. Which of the following statements about f are true?

- I. f is continuous at $x = 0$.
 II. f is differentiable at $x = 0$.
 III. f has an absolute minimum at $x = 0$.

- (A) I only (B) II only (C) III only (D) I and III only (E) II and III only

82. If f is a continuous function and if $F'(x) = f(x)$ for all real numbers x , then $\int_1^3 f(2x) dx =$

- (A) $2F(3) - 2F(1)$
 (B) $\frac{1}{2}F(3) - \frac{1}{2}F(1)$
 (C) $2F(6) - 2F(2)$
 (D) $F(6) - F(2)$
 (E) $\frac{1}{2}F(6) - \frac{1}{2}F(2)$

83. If $a \neq 0$, then $\lim_{x \rightarrow a} \frac{x^2 - a^2}{x^4 - a^4}$ is

- (A) $\frac{1}{a^2}$ (B) $\frac{1}{2a^2}$ (C) $\frac{1}{6a^2}$ (D) 0 (E) nonexistent


84. Population y grows according to the equation $\frac{dy}{dt} = ky$, where k is a constant and t is measured in years. If the population doubles every 10 years, then the value of k is
- (A) 0.069 (B) 0.200 (C) 0.301 (D) 3.322 (E) 5.000

x	2	5	7	8
$f(x)$	10	30	40	20

85. The function f is continuous on the closed interval $[2, 8]$ and has values that are given in the table above. Using the subintervals $[2, 5]$, $[5, 7]$, and $[7, 8]$, what is the trapezoidal approximation of

$$\int_2^8 f(x) dx ?$$

- (A) 110 (B) 130 (C) 160 (D) 190 (E) 210



86. The base of a solid is a region in the first quadrant bounded by the x -axis, the y -axis, and the line $x + 2y = 8$, as shown in the figure above. If cross sections of the solid perpendicular to the x -axis are semicircles, what is the volume of the solid?
- (A) 12.566 (B) 14.661 (C) 16.755 (D) 67.021 (E) 134.041

87. Which of the following is an equation of the line tangent to the graph of $f(x) = x^4 + 2x^2$ at the point where $f'(x) = 1$?

- (A) $y = 8x - 5$
 (B) $y = x + 7$
 (C) $y = x + 0.763$
 (D) $y = x - 0.122$
 (E) $y = x - 2.146$

88. Let $F(x)$ be an antiderivative of $\frac{(\ln x)^3}{x}$. If $F(1) = 0$, then $F(9) =$

- (A) 0.048 (B) 0.144 (C) 5.827 (D) 23.308 (E) 1,640.250

89. If g is a differentiable function such that $g(x) < 0$ for all real numbers x and if $f'(x) = (x^2 - 4)g(x)$, which of the following is true?

- (A) f has a relative maximum at $x = -2$ and a relative minimum at $x = 2$.
 (B) f has a relative minimum at $x = -2$ and a relative maximum at $x = 2$.
 (C) f has relative minima at $x = -2$ and at $x = 2$.
 (D) f has relative maxima at $x = -2$ and at $x = 2$.
 (E) It cannot be determined if f has any relative extrema.

90. If the base b of a triangle is increasing at a rate of 3 inches per minute while its height h is decreasing at a rate of 3 inches per minute, which of the following must be true about the area A of the triangle?
- (A) A is always increasing.
 (B) A is always decreasing.
 (C) A is decreasing only when $b < h$.
 (D) A is decreasing only when $b > h$.
 (E) A remains constant.
91. Let f be a function that is differentiable on the open interval $(1, 10)$. If $f(2) = -5$, $f(5) = 5$, and $f(9) = -5$, which of the following must be true?
- I. f has at least 2 zeros.
 II. The graph of f has at least one horizontal tangent.
 III. For some c , $2 < c < 5$, $f(c) = 3$.
- (A) None
 (B) I only
 (C) I and II only
 (D) I and III only
 (E) I, II and III
92. If $0 \leq k < \frac{\pi}{2}$ and the area under the curve $y = \cos x$ from $x = k$ to $x = \frac{\pi}{2}$ is 0.1, then $k =$
- (A) 1.471 (B) 1.414 (C) 1.277 (D) 1.120 (E) 0.436

ANS

- 76.A 77.C 78.B 79.A 80.B 81.D 82.E 83.B 84.A 85.C 86.C 87.D
 88.C 89.B 90.D 91.E 92.D

Part II

1. Let R be the region bounded by the x -axis, the graph of $y = \sqrt{x}$, and the line $x = 4$.
- (a) Find the area of the region R .
- (b) Find the value of h such that the vertical line $x = h$ divides the region R into two regions of equal area.
- (c) Find the volume of the solid generated when R is revolved about the x -axis.
- (d) The vertical line $x = k$ divides the region R into two regions such that when these two regions are revolved about the x -axis, they generate solids with equal volumes. Find the value of k .

(a) $\frac{16}{3}$ (b) $h = \sqrt[3]{16}$ or 2.520 (c) 8π or 25.133 (d) $k = \sqrt{8}$ or 2.828

2. Let f be the function given by $f(x) = 2xe^{2x}$.

(a) Find $\lim_{x \rightarrow -\infty} f(x)$ and $\lim_{x \rightarrow \infty} f(x)$.

(b) Find the absolute minimum value of f . Justify that your answer is an absolute minimum.

(c) What is the range of f ?

(d) Consider the family of functions defined by $y = bxe^{bx}$, where b is a nonzero constant. Show that the absolute minimum value of bxe^{bx} is the same for all nonzero values of b .


(a) $\lim_{x \rightarrow -\infty} 2xe^{2x} = 0$, $\lim_{x \rightarrow \infty} 2xe^{2x} = \infty$ (b) $-\frac{1}{e}$ (c) Range of $f = [-\frac{1}{e}, \infty)$

(d) $y' = be^{bx} + b^2xe^{bx} = be^{bx}(1 + bx) = 0$

if $x = -1/b$

At $x = -1/b$, $y = -1/e$

y has an absolute minimum value of $-1/e$ for all nonzero b



t (seconds)	$v(t)$ (feet per second)
0	0
5	12
10	20
15	30
20	55
25	70
30	78
35	81
40	75
45	60
50	72

3. The graph of the velocity $v(t)$, in ft/sec, of a car traveling on a straight road, for $0 \leq t \leq 50$, is shown above. A table of values for $v(t)$, at 5 second intervals of time t , is shown to the right of the graph.

- (a) During what intervals of time is the acceleration of the car positive? Give a reason for your answer.
- (b) Find the average acceleration of the car, in ft/sec^2 , over the interval $0 \leq t \leq 50$.
- (c) Find one approximation for the acceleration of the car, in ft/sec^2 , at $t = 40$. Show the computations you used to arrive at your answer.
- (d) Approximate $\int_0^{50} v(t) dt$ with a Riemann sum, using the midpoints of five subintervals of equal length. Using correct units, explain the meaning of this integral.

(a) Acceleration is positive on $(0, 35)$ and $(45, 50)$ because the velocity $v(t)$ is increasing on $[0, 35]$ and $[45, 50]$

(b) $\frac{72}{50}$

(c) Difference quotient; e.g.

$$\frac{v(45) - v(40)}{5} = \frac{60 - 75}{5} = -3 \text{ ft}/\text{sec}^2 \text{ or}$$

$$\frac{v(40) - v(35)}{5} = \frac{75 - 81}{5} = -\frac{6}{5} \text{ ft}/\text{sec}^2 \text{ or}$$

$$\frac{v(45) - v(35)}{10} = \frac{60 - 81}{10} = -\frac{21}{10} \text{ ft}/\text{sec}^2$$

(d) $\int_0^{50} v(t) dt$

$$\begin{aligned} &\approx 10[v(5) + v(15) + v(25) + v(35) + v(45)] \\ &= 10(12 + 30 + 70 + 81 + 60) \\ &= 2530 \text{ feet} \end{aligned}$$

This integral is the total distance traveled in feet over the time 0 to 50 seconds.

4. Let f be a function with $f(1) = 4$ such that for all points (x, y) on the graph of f the slope is given by $\frac{3x^2 + 1}{2y}$.

(a) Find the slope of the graph of f at the point where $x = 1$.

(b) Write an equation for the line tangent to the graph of f at $x = 1$ and use it to approximate $f(1.2)$.

(c) Find $f(x)$ by solving the separable differential equation $\frac{dy}{dx} = \frac{3x^2 + 1}{2y}$ with the initial condition $f(1) = 4$.

(d) Use your solution from part (c) to find $f(1.2)$.


(a) $\frac{1}{2}$ (b) $f(1.2) \approx 4.1$ (c) $f(x) = \sqrt{x^3 + x + 14}$ (d) $f(1.2) \approx 4.114$

5. The temperature outside a house during a 24-hour period is given by

$$F(t) = 80 - 10 \cos\left(\frac{\pi t}{12}\right), \quad 0 \leq t \leq 24,$$

where $F(t)$ is measured in degrees Fahrenheit and t is measured in hours.

(a) Sketch the graph of F on the grid below.



(b) Find the average temperature, to the nearest degree Fahrenheit, between $t = 6$ and $t = 14$.

- (c) An air conditioner cooled the house whenever the outside temperature was at or above 78 degrees Fahrenheit. For what values of t was the air conditioner cooling the house?
- (d) The cost of cooling the house accumulates at the rate of \$0.05 per hour for each degree the outside temperature exceeds 78 degrees Fahrenheit. What was the total cost, to the nearest cent, to cool the house for this 24-hour period?

(a) (b) $\approx 87^{\circ}F$ (c) $5.230 \leq t \leq 18.769$ (d) ≈ 5.10

6. Consider the curve defined by $2y^3 + 6x^2y - 12x^2 + 6y = 1$.

(a) Show that $\frac{dy}{dx} = \frac{4x - 2xy}{x^2 + y^2 + 1}$.

(b) Write an equation of each horizontal tangent line to the curve.

(c) The line through the origin with slope -1 is tangent to the curve at point P . Find the x - and y -coordinates of point P .

(a)

(b) $y=0.165$

(c) $x = -\frac{1}{2}, y = \frac{1}{2}$